

Hemiklonale Wasserfrösche – Hemiclonal water frogs

- 2018 Dolezalkova-Kastankova M, Pruvost NBM, Plötner J, **Reyer H-U**, Janko K, Choleva L (2018): All-male hybrids of a tetrapod *Pelophylax esculentus* share its origin and genetics of maintenance. *Biol. Sex Differ.* 9: Article 13.
- 2015 Hoffmann A, Abt Tietje G, **Reyer H-U** (2015): Spatial behavior in relation to mating systems: movement patterns, nearest neighbour distances, and mating success in diploid and polyploid frog hybrids (*Pelophylax esculentus*). *Behav. Ecol. Sociobiol.* 69: 501-517.
- Hoffmann A, Plötner J, Pruvost NBM, Christiansen DG, Röthlisberger S, Choleva L, Mikulíček P, Cogălniceanu D, Sas- Kovács I, Shabanov D, Morozov-Leonov S, **Reyer H-U** (2015): Genetic diversity and distribution patterns of diploid and polyploid hybrid water frog populations (*Pelophylax esculentus* complex) across Europe. *Mol. Ecol.* 24: 4371-4391.
- Pruvost NBM, Mikulicek P, Choleva L, **Reyer H-U** (2015): Contrasting reproductive strategies of triploid hybrid males in vertebrate mating systems. *J. Evol. Biol.* 28: 189-204.
- Reyer H-U**, Arioli-Jakob C, Arioli M (2015): Post-zygotic selection against parental genotypes during larval development maintains all-hybrid populations of the frog *Pelophylax esculentus*. *BMC Evol. Biol.* 15: 131.
- 2013 Hoffmann A, **Reyer H-U** (2013): Genomic effects on advertisement call structure in diploid and triploid hybrid water frogs (*Anura, Pelophylax esculentus*). *BMC Ecol.* 13: 47.
- Pruvost NBM, Hoffmann A, **Reyer H-U** (2013): Gamete production patterns, ploidy, and population genetics reveal evolutionary significant units in hybrid water frogs (*Pelophylax esculentus*). *Ecol. Evol.* 3: 2933-2946.
- Pruvost NBM, Hollinger O & **Reyer H-U** (2013): Genotype-temperature interactions on larval performance shape population structure in hybridogenetic water frogs (*Pelophylax esculentus* complex). *Funct. Ecol.* 27: 459-471.
- Stohr AC, Hoffmann A, Papp T, Robert N, Pruvost NBM, **Reyer H-U** & Marschang RE (2013): Long-term study of an infection with Rana viruses in a group of edible frogs (*Pelophylax kl. esculentus*) and partial characterization of two viruses based on four genomic regions. *Vet. J.* 197:238-244.
- 2012 Embrechts E & **Reyer H-U** (2012): Age and size of hybrid water frogs: the role of genotype and ecology. *Herpetologica* 68: 468-481.
- 2011 Christiansen DG & **Reyer H-U** (2011): effects of geographic distance, sea barriers and habitat on the genetic structure and diversity of all-hybrid water frog populations. *Heredity* 106: 25-36.2009
- 2009 Christiansen DG & **Reyer H-U** (2009): From clonal to sexual hybrids: genetic recombination via triploids in all-hybrid populations of water frogs. *Evolution* 63: 1754-1768.
- Vorburger C, Schmelter OS, Hotz H, Guex G-O & **Reyer H-U** (2009): Masked damage: mutational load in hemiclonal water frogs. In: Van Oijk P., Martens K. Schoen I. (eds): *Lost Sex -- The Evolutionary Biology of Parthenogenesis*; pp. 433-446. Springer Press, Berlin.

- 2008 Plötner J, Beerli P, Uzzell T, Spolsky C, Ohst T, Litvinchuk SN, Guex G-O, **Reyer H-U** & Hotz H (2008): Widespread unidirectional transfer of mitochondrial DNA: a case in western Palearctic water frogs. *J. Evol. Biol.* 21: 668-681.
- Reyer, HU** (2008): Mating with the wrong species can be right. *Trends Ecol. Evol.* 23: 289-292.
- 2007 Som C, Bagheri HC & **Reyer H-U** (2007): Mutation accumulation and fitness effects in hybridogenetic populations: a comparison to sexual and asexual systems. *BMC Evol. Biol.* 7: 80.
- Waelti MO & **Reyer H-U** (2007): Food supply modifies the trade-off between past and future reproduction in a sexual parasite-host system (*Rana esculenta*, *R. lessonae*). *Oecologia* 152: 415-424.
- Som C & **Reyer H-U** (2007): Hemiclonal reproduction slows down the speed of Muller's ratchet in the hybridogenetic frog, *Rana esculenta*. *J. Evol. Biol.* 20: 650-660.
- 2006 Som C & **Reyer H-U** (2006a): Variation in sex ratio and evolutionary rate of hemiclonal *Rana esculenta* populations. *Evol. Ecol.* 20: 159-17.
- Som C & **Reyer H-U** (2006b): Demography and evolution of pure diploid-triploid water frog (*Rana esculenta*) populations. *Evol. Ecol. Res.* 8: 1235-1248. •
- 2004 Abt Tietje G & **Reyer H-U** (2004): Larval development and recruitment of juveniles in a natural population of *Rana lessonae* and *Rana esculenta*. *Copeia* 3: 638-646.
- Reyer H-U** & Bättig I (2004): Identification of reproductive status in female frogs - A quantitative comparison of nine methods. *Herpetologica* 60: 349-357.
- Reyer H-U**, Waelti MO, Bättig I, Altwegg R & Hellriegel B (2004): Low proportions of reproducing hemiclonal females increase the stability of a sexual parasite-host system (*Rana esculenta*, *Rana lessonae*). *J. Anim. Ecol.* 73: 1089-1101.
- 2003 **Reyer H-U**, Niederer B & Hettyey A. (2003): Variation in fertilisation abilities between hemiclonal hybrid and sexual parental males of sympatric water frogs (*Rana lessonae*, *R. esculenta*, *R. ridibunda*). *Behav. Ecol. Sociobiol.* 54: 274-284.
- Altwegg R & **Reyer H-U** (2003): Fitness consequences of larval growth environment and timing of metamorphosis in water frogs. *Evolution* 57: 872-882.
- Vorburger C & **Reyer H-U** (2003): A genetic mechanism of species replacement in European water frogs? *Conserv. Gen.* 4: 141-155.
- 2002 Holenweg Peter A-K, **Reyer H-U** & Abt Tietje G (2002): Species and sex ratio differences in mixed populations of hybridogenetic water frogs: the influence of pond features. *Ecoscience* 9: 1-11.
- 2001 Holenweg Peter A-K, **Reyer H-U** & Abt Tietje G (2001): Homing behavior of *Rana lessonae*, *R. ridibunda* and their hybridogenetic associate *R. esculenta* after experimental displacement. *Amphibia-Reptilia* 22: 475-480.
- Engeler B & **Reyer H-U** (2001): Choosy females and indifferent males: mate choice in mixed populations of the water frogs *Rana lessonae* and *Rana esculenta*. *Behav. Ecol.* 12: 600-606.
- Negovetic S, Anholt BA, Semlitsch RD & **Reyer H-U** (2001): Specific responses of sexual and hybridogenetic European water frog tadpoles to temperature. *Ecology* 82: 766-774.

- Thurnheer S & **Reyer H-U** (2001): Spatial distribution and survival rate of water frog tadpoles in relation to biotic and abiotic factors. A field experiment. *Amphibia-Reptilia* 22: 21-32.
- 2000 Garner TWJ, Gautschi B, Rothlisberger S & **Reyer H-U** (2000): A set of A repeat micro-satellite markers derived from the pool frog, *Rana lessonae*. *Mol. Ecol.* 9: 2173-2175.
- Hellriegel B & **Reyer H-U** (2000): Factors influencing the composition of mixed populations of a hemiclinal hybrid and its sexual host. *J. Evol. Biol.* 13: 906-918.
- Holenweg A-K & **Reyer H-U** (2000): Hibernating behavior of *Rana lessonae* and *R. esculenta* in their natural habitat. *Oecologia* 123: 41-47.
- Roesli M & **Reyer H-U** (2000): Male vocalisation and female choice in the hybridogenetic *Rana lessonae/R. esculenta* complex. *Anim. Behav.* 60: 745-755.
- Som C, Anholt BR & **Reyer H-U** (2000): The effect of assortative mating on the coexistence of a hybridogenetic water frog and its sexual host. *Am. Nat.* 156: 34-46.
- 1999 **Reyer H-U**, Frei G & Som C (1999): Cryptic female choice: frogs reduce clutch size when amplexed by undesired males. *Proc. R. Soc. Lond. B* 266: 2101-2107.
- 1997 Bergen K, Semlitsch RD & **Reyer H-U** (1997): Hybrid female matings are directly related to the availability of *Rana lessonae* and *Rana esculenta* males in experimental populations. *Copeia* 1997: 275-283.
- Fioramonti E, Semlitsch RD, **Reyer H-U** & Fent K (1997): Effects of tri-phenyltin and pH on the growth and development of tadpoles of *Rana lessonae* and *Rana esculenta*. *Environm. Toxicol. Chem.* 16: 1940-1947.
- Rist L, Semlitsch RD, Hotz H & **Reyer H-U** (1997): Feeding behaviour, food consumption, and growth efficiency of tadpoles of the *Rana esculenta* complex. *Funct. Ecol.* 11: 735-742.
- 1993 Abt G & **Reyer H-U** (1993): Mate choice and fitness in a hybrid frog: *Rana esculenta* females prefer *Rana lessonae* males over their own. *Behav. Ecol. Sociobiol.* 32: 221-228.
- 1992 Semlitsch RD & **Reyer H-U** (1992a): Modification of anti-predator behaviour in tadpoles by environmental conditioning. *J. Anim. Ecol.* 61: 353-360.
- Semlitsch RD & **Reyer H-U** (1992b): Performance of tadpoles from the hybridogenetic *Rana esculenta* complex: Interactions with pond drying and interspecific competition. *Evolution* 46: 665-676.